


Retrained generic antibodies can recognize SARS-CoV-2

3 February 2021, by Brian Flood



Double-faced peptide-based boosters are computationally designed to allow recognition of SARS-CoV-2 (grey, schematic) by Hepatitis B antibodies. One booster face made of ACE2-mimic peptides (red) can bind to the receptor binding domain of SARS-CoV-2 (blue). The other booster face composed of a Hepatitis B core-antigen (orange) can target the Hepatitis B antibody fragment (green). This approach can provide a cheap and efficient neutralization of emerging pathogens by generic antibodies. Credit: University of Illinois at Chicago

SARS-CoV-2, the new coronavirus behind the current pandemic, infects humans by binding its surface-exposed spike proteins to ACE2 receptors exposed on the cell membranes.

Upon a vaccination or a real infection, it takes several weeks before the immunity develops antibodies that can selectively bind to these spike proteins. Such antibody-labeled viruses are neutralized by the natural killer and T cells operated by the human immunity.

An alternative approach to train the immunity response is offered by researchers at the University of Illinois Chicago and California State University at Sacramento who have developed a novel strategy that redirects antibodies for other diseases existing in humans to the spike proteins of SARS-CoV-2.

In their study published by the *Journal of Physical Chemistry Letters*, the team proposes using peptide-based double-faced "booster" inhibitors, with one face binding to the spike proteins of SARS-CoV-2 and the other face binding to generic hepatitis B antibodies.

"Once the SARS-CoV-2 viruses become labeled by the hepatitis B antibodies via intermediate boosters, the viruses will be neutralized. This universal approach allows a dramatic shortening of the response time upon real infections, which can be critical in certain patients or conditions," said Petr Král, UIC professor of chemistry, physics, [pharmaceutical sciences](#) and chemical engineering, and senior author on the paper.

Král and Yanxiao Han, who recently earned a Ph.D. in chemistry at UIC and is first author on the paper, believe the study could provide guidance in the preparation of generic therapeutics against emerging pathogens with the combined advantages of small-[protein](#) and antibody therapies.

"The dramatic impact which novel viruses can have on humans could be fast mitigated in the absence of their vaccination if generic [antibodies](#) present within them are temporarily retrained to recognize these viruses," the researchers wrote.

In a [study published last spring](#), Král and Han extracted different peptides from ACE2 that interact directly with the viral spike protein.

"We investigated potential COVID-19 therapeutics using [computer simulations](#) based on the X-ray crystal structure of the receptor-binding domain of SARS-CoV-2 when it is bound to ACE2," Král said. "Similar to our latest study, identifying these kinds of inhibitors could lead to new treatments to combat the coronavirus."

More information: Yanxiao Han et al. Retrained Generic Antibodies Can Recognize SARS-CoV-2, *The Journal of Physical Chemistry Letters* (2021). [DOI: 10.1021/acs.jpcllett.0c03615](https://doi.org/10.1021/acs.jpcllett.0c03615)

Provided by University of Illinois at Chicago

APA citation: Retrained generic antibodies can recognize SARS-CoV-2 (2021, February 3) retrieved 28 May 2022 from <https://medicalxpress.com/news/2021-02-retrained-antibodies-sars-cov-.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.