

# SARS-CoV-2 antibodies detected six months after infection

9 February 2021



were seropositive in month 1 and had reported symptoms or had a positive polymerase chain reaction test. Of these 705 participants, 12.2 percent sero-reverted within six months of symptom onset. Only five participants (

(HealthDay)—For many individuals, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) antispikes remain for at least six months after infection, according to a report published by the U.K. Biobank.

To determine the extent of past infection with SARS-CoV-2 in different population subgroups, the U.K. Biobank collected [blood samples](#) from about 20,200 individuals on a monthly basis for six months.

A total of 18,893 individuals (93.5 percent) provided at least one sample that was successfully assayed between May 27 and Dec. 4, 2020. The researchers found that 9 percent of individuals had at least one sample that was seropositive for SARS-CoV-2 antispikes during the study period. Seroprevalence differed significantly by age, ethnicity, geographic region, and [socioeconomic status](#). Overall, 1,264 individuals were seropositive in month 1. The duration of seropositivity was assessed among those who

APA citation: SARS-CoV-2 antibodies detected six months after infection (2021, February 9) retrieved 12 October 2022 from <https://medicalxpress.com/news/2021-02-sars-cov-antibodies-months-infection.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*