


The important role of music in neurorehabilitation: Filling in critical gaps

March 10 2021



Regions of interest (ROIs) identified around the therapist and child in A) Music-Based Intervention and B) Non-Music Control Intervention. Credit: Faculty of Music and Faculty of Medicine, University of Toronto

Music-based interventions have become a core ingredient of effective neurorehabilitation in the past 20 years thanks to the growing body of knowledge. In this theme issue of *Neurorehabilitation*, experts in the field highlight some of the current critical gaps in clinical applications that have been less thoroughly investigated, such as post-stroke cognition, traumatic brain injury, and autism and specific learning disabilities.

Neurologic Music Therapy is the clinical and evidence-based use of [music](#) interventions by a credentialed professional. Research in the 1990s showed for the first time how musical-rhythmic stimuli can improve mobility in stroke and Parkinson's disease patients. We now know that music-based interventions can effectively address a wide range of impairments in sensorimotor, speech/language, and cognitive functions.

"The use of music-based interventions in neurorehabilitation was virtually unknown 25 years ago," explained Guest Editor Michael Thaut, Ph.D., Director, Music and Health Science Research Collaboratory, Faculty of Music and Faculty of Medicine, University of Toronto. "Since then, a growing body of research has shown how musical-rhythmic stimuli can improve mobility disorders such as stroke and Parkinson's disease, and music-based interventions have now become a core ingredient of effective neurorehabilitation. For example, Rhythmic Auditory Stimulation (RAS) has now been adopted in several official stroke care guidelines in the United States and Canada."

This collection of articles includes three studies on the use of music in [traumatic brain injury](#) rehabilitation; two studies looking at music-based interventions in children with autism and learning disabilities, respectively; the little-investigated connection between motor training

and cognitive outcomes in chronic stroke rehabilitation; and a theoretical paper on mechanisms of neuroplastic changes underlying successful Neurologic Music Therapy interventions that provides a theoretical understanding of how music shapes brain function in neurorehabilitation on an impairment level. Several papers in the issue review research into the treatment system of Neurologic Music Therapy that has been endorsed by the World Federation of Neurorehabilitation as evidence-based and is practiced by certified clinicians in over 50 countries.

Lead investigator Catherine M. Haire, Ph.D., Faculty of Music, Music and Health Science Research Collaboratory, University of Toronto, and colleagues report on results of a randomized controlled trial of therapeutic instrumental music performance (TIMP) with and without motor imagery on chronic post-stroke cognition and affect. They found that the mental flexibility aspect of executive functioning appears to be enhanced by therapeutic instrumental music training in conjunction with motor imagery, possibly due to multisensory integration and consolidation of representations through motor imagery rehearsal following active practice. "Active training using musical instruments appears to have a positive impact on affective responding," commented Dr. Haire. "However, these changes occurred independently of improvements to cognition."

The effectiveness of music-based interventions in autism has been recognized for decades, but there has been little empirical investigation of the processes involved and how they compare to other approaches. Aparna Nadig, Ph.D., School of Communication Sciences and Disorders, McGill University, and colleagues found that compared to a non-music control [intervention](#), children in music-based interventions spent more time engaged in triadic engagement (between child, therapist, and activity) and produced greater movement, depending on the type of musical instrument involved. "Taken together, these findings provide helpful initial evidence of the active ingredients of music-based

interventions in autism," noted Dr. Nadig.

Looking ahead, Dr. Thaut commented that "A significant trend is the move from a therapy approach to a learning/training approach that allows the patient to become a more autonomous and independent participant in therapy. Providing patients with music-based devices for more independent and more frequent training via music technology will be an important new development. Future challenges will be to develop approaches and build technology to integrate Neurologic Music Therapy into telehealth post-COVID-19 to reach more patients in need worldwide who do not have access to widely distributed neurorehabilitation services. We are at a point where we can state clinically that the brain that engages in music, is changed by engaging in music."

More information: Catherine M. Haire et al, Effects of therapeutic instrumental music performance and motor imagery on chronic post-stroke cognition and affect: A randomized controlled trial, *NeuroRehabilitation* (2021). [DOI: 10.3233/NRE-208014](https://doi.org/10.3233/NRE-208014)

Provided by IOS Press

Citation: The important role of music in neurorehabilitation: Filling in critical gaps (2021, March 10) retrieved 30 March 2023 from <https://medicalxpress.com/news/2021-03-important-role-music-neurorehabilitation-critical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.