

Risk of death for men 60% higher than for women in study of 28 countries

15 March 2021



Credit: Unsplash/CC0 Public Domain

A large study of people in 28 countries found men aged 50 and over had a 60% greater risk of death than women, partly explained by heavier rates of smoking and heart disease in men, although the gap varied across countries, according to new research in *CMAJ (Canadian Medical Association Journal)*.

"Many studies have examined the potential impact of social, behavioral and [biological factors](#) on [sex differences](#) in mortality, but few have been able to investigate potential variation across countries," writes Dr. Yu-Tzu Wu, Institute of Psychiatry, Psychology and Neuroscience, King's College London, and Population Health Sciences Institute, Newcastle University, United Kingdom, with coauthors. "Different cultural traditions, historical contexts, and economic and societal development may influence gender experiences in different countries, and thus variably affect the health status of men and women."

The study examined different socioeconomic (education, wealth), lifestyle (smoking, [alcohol consumption](#)), health (heart diseases, diabetes,

hypertension and depression) and social (spouse, living alone) factors that might contribute to the mortality gap between men and women aged 50 and older. The data included more than 179,000 people across 28 countries and more than half (55%) were women.

"[T]he effects of sex on mortality should include not only physiologic variation between men and women but also the social construct of gender, which differs across societies. In particular, the large variation across countries may imply a greater effect of gender than sex. Although the biology of the sexes is consistent across populations, variation in cultural, societal and historical contexts can lead to different life experiences of men and [women](#) and variation in the mortality gap across countries."

The findings are consistent with the literature on life expectancy and death rates.

"The heterogeneity of sex differences in mortality across countries may indicate the substantial impact of gender on healthy aging in addition to biological sex, and the crucial contributions of smoking may also vary across different populations," write the authors.

The researchers recommend that public health policies should account for sex- and gender-based differences and the influence of social and [cultural factors](#) on health.

More information: "Sex differences in mortality: results from a population-based study of 12 longitudinal cohorts" is published March 15, 2021. www.cmaj.ca/lookup/doi/10.1503/cmaj.200484

Provided by Canadian Medical Association Journal

APA citation: Risk of death for men 60% higher than for women in study of 28 countries (2021, March 15) retrieved 25 April 2021 from <https://medicalxpress.com/news/2021-03-death-men-higher-women-countries.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.