


Prostate cancer uses metabolic switch to thrive after hormone therapy

22 March 2021



Micrograph showing prostatic acinar adenocarcinoma (the most common form of prostate cancer) Credit: Wikipedia

Studying the cellular metabolism of prostate cancer, a team of Duke Health-led researchers identified a key reason hormone therapies eventually fail, while also laying out a way to bypass the problem using an entirely new therapeutic approach.

The findings, published during the week of March 22 in the Proceedings of the National Academy of Sciences, describe how hormone therapies target the androgen receptor to essentially starve tumor cells of a crucial fuel source. This initially works well to halt tumor growth, but then the cancer cells compensate, switching to a different enzyme to exploit the fuel and proliferating as they become resistant to hormone therapies.

The team of Duke Cancer Institute researchers used that finding to propose a treatment strategy that eliminates the need to inhibit the androgen receptor completely. Their goal is to directly target the tumor's preferred source of fuel—an amino acid "Our work demonstrates this metabolic switch to be called glutamine.

In studies using prostate cancer cell lines, human prostate cancer tissue and animal models, the novel therapeutic strategy successfully inhibited tumor growth. Clinical trials are in being planned using a currently available drug that inhibits glutamine use by tumor cells.

"Instead of inhibiting androgen receptor using hormonal therapy, a better therapeutic strategy is to inhibit glutamine utilization directly," said senior author Jiaoti Huang, M.D., Ph.D., chair of Duke's Department of Pathology.

"Since glutamine is not essential for normal tissue, there will be fewer side effects, which is one of the biggest downsides to hormonal therapies," Huang said. "Direct inhibition of the enzyme that controls glutamine utilization would also make it more difficult for tumor cells to develop resistance."

Huang and co-authors—including Daniel George, M.D., professor in the departments of Medicine and Surgery at Duke who leads the clinical trial design—initiated the study to better understand prostate cancer cell metabolism, which still has many unknowns.

They found that hormone therapy initially inhibits a certain form of glutamine-converting enzyme called kidney-type glutaminase (KGA). This KGA enzyme depends on the androgen receptor and makes it possible for cancer cells to use glutamine. By suppressing it, hormonal therapies successfully slow cancer growth for a time.

But the tumor cells eventually find a workaround, switching to a different enzyme—glutaminase C (GAC)—which doesn't rely on androgen receptor. When tumors make this switch to GAC, they proliferate aggressively, becoming castrationresistant prostate cancer.

one of the key mechanisms in therapeutic



resistance and disease progression," George said.

By targeting glutamine metabolism, the researchers pioneered a way to bypass the complex <u>androgen</u> <u>receptor</u> signaling processes, instead directly suppressing the production of energy and building blocks required by prostate cancer cells, essentially starving <u>tumor cells</u> to death.

"Since metabolic activity directly controls cellular proliferation, it may be more difficult for the <u>tumor cells</u> to overcome a metabolic inhibition to develop resistance," Huang said. "Our study shows that pharmacological inhibition of GAC can significantly suppress castration-resistant prostate cancer."

More information: Lingfan Xu el al., "A glutaminase isoform switch drives therapeutic resistance and disease progression of prostate cancer," *PNAS* (2021). www.pnas.org/cgi/doi/10.1073/pnas.2012748118

Provided by Duke University Medical Center APA citation: Prostate cancer uses metabolic switch to thrive after hormone therapy (2021, March 22) retrieved 19 April 2021 from

https://medicalxpress.com/news/2021-03-prostate-cancer-metabolic-hormone-therapy.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.