

Researchers urge a closer examination of sunlight's efficacy in inactivating the SARS-CoV-2 virus

31 March 2021, by Sonia Fernandez



Credit: CC0 Public Domain

A year ago scientists everywhere were scrambling to get their minds around the SARS-CoV-2, a novel coronavirus that caused the pandemic from which we are only now beginning to emerge. The world clung to every new development, every bit of science that could provide clues to managing life in the presence of this mysterious killer.

Many science-backed COVID-19 management concepts remain unchanged to this day: handwashing with soap and warm water disrupts the virus' lipid membrane. Social distancing can attenuate the virus's spread, ideally keeping it out of a host until it degrades. Other notions, such as droplet contact being the primary mode of transmission, were modified when emerging evidence showed that under certain conditions, the virus could remain suspended in air for extended periods of time.

In a [letter](#) in the *Journal of Infectious Diseases*, a team of researchers from UC Santa Barbara, Oregon State University, University of Manchester

and ETH Zurich examines another of SARS-CoV-2's well known characteristics—its vulnerability to sunlight. Their conclusion? It might take more than UV-B rays to explain sunlight inactivation of SARS-CoV-2.

The idea that an additional mechanism might be in play came when the team compared data from a July 2020 [study](#) that reported rapid sunlight inactivation of SARS-CoV-2 in a lab setting, with a [theory](#) of coronavirus inactivation by [solar radiation](#) that was published just a month earlier.

"The theory assumes that inactivation works by having UV-B hit the RNA of the virus, damaging it," said UC Santa Barbara mechanical engineering professor and lead author Paolo Luzzatto-Fegiz. Judging from the discrepancies between the experimental results and the predictions of the theoretical model, however, the research team felt that RNA inactivation by UV-B "might not be the whole story."

According to the letter, the experiments demonstrated virus inactivation times of about 10-20 minutes—much faster than predicted by the theory.

"The theory predicts that inactivation should happen an order of magnitude slower," Luzzatto-Fegiz said. In the experiments, viruses in simulated saliva and exposed to UV-B lamps were inactivated more than eight times faster than would have been predicted by the theory, while those cultured in a complete growth medium before exposure to UV-B were inactivated more than three times faster than expected. To make the math of the theory fit the data, according to the letter, SARS-CoV-2 would have to exceed the highest UV-B sensitivity of any currently known virus.

Or, Luzzato-Fegiz and colleagues reasoned, there could be another mechanism at play aside from RNA inactivation by UV-B rays. For instance, UV-A, another, less energetic component of sunlight might be playing a more active role than previously thought.

"People think of UV-A as not having much of an effect, but it might be interacting with some of the molecules in the medium," he said. Those reactive intermediate molecules in turn could be interacting with the virus, hastening inactivation. It's a concept familiar to those who work in [wastewater treatment](#) and other environmental science fields.

"So, scientists don't yet know what's going on," Luzzato-Fegiz said; "Our analysis points to the need for additional experiments to separately test the effects of specific light wavelengths and medium composition."

Results of such experiments might provide clues into new ways of managing the virus with widely available and accessible UV-A and UV-B radiation. While UV-C radiation is proved effective against SARS-CoV-2, this wavelength does not reach the earth's surface and must be manufactured. Although UV-C is presently used in air filtration and in other settings, its short wavelengths and [high energy](#) also makes UV-C the most damaging form of UV radiation, limiting its practical application and raising other safety concerns.

"UV-C is great for hospitals," said co-author Julie McMurry. "But in other environments—for instance kitchens or subways—UV-C would interact with the particulates to produce harmful ozone." While no single intervention will eliminate risk, this research would provide one further tool to reduce exposure, thus slowing transmission and improving health outcomes.

Co-author and UCSB mechanical engineering professor Yangying Zhu added that UV-A turning out to be capable of inactivating the [virus](#) could be very advantageous: there are now widely available inexpensive LED bulbs that are many times stronger than natural sunlight, which could accelerate inactivation times. UV-A could potentially be used far more broadly to augment air

filtration systems at relatively low risk for human health, especially in high-risk settings such as hospitals and public transportation, but the specifics of each setting warrant consideration, said co-author Fernando Temprano-Coleto.

More information: Paolo Luzzato-Fegiz et al. UVB Radiation Alone May Not Explain Sunlight Inactivation of SARS-CoV-2, *The Journal of Infectious Diseases* (2021). [DOI: 10.1093/infdis/jiab070](#)

Provided by University of California - Santa Barbara

APA citation: Researchers urge a closer examination of sunlight's efficacy in inactivating the SARS-CoV-2 virus (2021, March 31) retrieved 11 October 2022 from <https://medicalxpress.com/news/2021-03-urge-closer-sunlight-efficacy-inactivating.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.