


COVID-19 in our dust may help predict outbreaks, study finds

13 April 2021, by Laura Arenschild



Credit: Pixabay/CC0 Public Domain

A study done in rooms where COVID-19 patients were isolated shows that the virus's RNA—part of the genetic material inside a virus—can persist up to a month in dust.

The study did not evaluate whether dust can transmit the [virus](#) to humans. It could, however, offer another option for monitoring COVID-19 outbreaks in specific buildings, including nursing homes, offices or schools.

Karen Dannemiller, senior author of the study, has experience studying dust and its relationship to potential hazards like mold and microbes.

"When the pandemic started, we really wanted to find a way that we could help contribute knowledge that might help mitigate this crisis," said Dannemiller, assistant professor of civil, environmental and geodetic engineering and environmental health sciences at The Ohio State University.

"And we've spent so much time studying dust and flooring that we knew how to test it."

The study, published today (April 13, 2021) in the journal *mSystems*, found some of the genetic material at the heart of the virus persists in dust, even though it is likely that the envelope around the virus may break down over time in dust. The envelope—the crown-like spiked sphere that contains the virus's material—plays an important role in the virus's transmission to humans.

The study offers another non-invasive avenue for monitoring buildings for COVID-19 outbreaks, especially as more people are vaccinated and return to communal spaces.

Municipalities and others have tested wastewater to evaluate the prevalence of COVID-19 in a given community—gene copies and fragments of the virus live in human waste, and by testing wastewater, local governments and others can determine how widespread the virus might be, even if people are asymptomatic.

Dust monitoring could offer similar understanding on a smaller scale—say, a specific nursing home, hospital or school.

"In nursing homes, for example, you're still going to need to know how COVID is spreading inside the building," said Nicole Renninger, lead author of the paper and an engineering graduate student in Dannemiller's lab. "For surveillance purposes, you need to know if you are picking up an outbreak that's going on right now."

For this study, the research team worked with the crews responsible for cleaning the rooms at Ohio State where students who tested positive for COVID-19 were isolated. They also collected samples from two homes where people who tested positive for COVID-19 lived. They gathered vacuum bags of dust from the cleaning crews and from the homes.

The researchers also tested swabs collected from

surfaces in the rooms.

They found genetic material from the SARS-CoV-2 virus—the virus that causes COVID-19—in 97% of the bulk dust samples and in 55% of the surface swabs.

The cleaning crews sprayed a chlorine-based disinfectant in the rooms prior to cleaning; the researchers believe that disinfectant destroyed the envelope and/or capsid—the outer coat surrounding the virus—likely defanging it for transmission.

The research team tested the samples when they arrived at the lab, shortly after the rooms were cleaned, then tested the samples again weekly. After four weeks, the virus's RNA did not significantly decay in the vacuum bags.

"We weren't sure that the [genetic material](#) would survive—there are many different organisms in dust, and we weren't sure we'd see any viral RNA at all," Renninger said. "And we were surprised when we found that the actual RNA itself seems to be lasting a pretty long time."

Testing dust to monitor for COVID-19 outbreaks would likely be most useful for smaller-scale communities with a high-risk population—a nursing home, for example, Dannemiller said. Testing indoor dust is also likely less expensive at that scale than testing wastewater or all individuals directly on a routine basis.

"We wanted to demonstrate that dust could be complementary to wastewater for surveillance," Dannemiller said. "Wastewater is great for a large population, but not everybody sheds the virus in feces, and you have to collect wastewater samples, which not everyone wants to do. People are already vacuuming these rooms, so [dust](#) may be a good option for some groups."

Even before this study was published, Dannemiller said the researchers presented their findings to an industry group that represents maintenance and cleaning staff, with a recommendation: "If they can wait at least an hour or more after a person leaves the room to clean, they should," she said, citing previous studies of viral viability on other materials

and in aerosols.

More information: Nicole Renninger et al. Indoor Dust as a Matrix for Surveillance of COVID-19. *mSystems* DOI: [10.1128/mSystems.01350-20](https://doi.org/10.1128/mSystems.01350-20)

Provided by The Ohio State University

APA citation: COVID-19 in our dust may help predict outbreaks, study finds (2021, April 13) retrieved 5 June 2022 from <https://medicalxpress.com/news/2021-04-covid-outbreaks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.