

Viewing your own face, even subconsciously, is rewarding

19 April 2021




Figure 1: Subliminal presentation of face (upper) and brain regions showing self-face related activation (bottom). Credit: Osaka University

As humans, we each have a powerful ability to easily recognize our own face. But now, researchers from Japan have uncovered new information about how our cognitive systems enable us to distinguish our own face from those of others, even when the information is presented subliminally.

In a study published this month in *Cerebral Cortex*, researchers from Osaka University have revealed that a central element of the dopamine reward pathway in the brain was activated when participants were subliminally shown images of their face. This provides new clues regarding the underlying processes of the brain involved in self-facial recognition.

When we are exposed to a subliminal image of our face—meaning we are not fully aware of it—many [brain regions](#) are activated in addition to those that process face information. Furthermore, our brain responds differently to supraliminal (conscious) and subliminal (subconscious) images of our face compared to [faces](#) of others. However, whether we use the same or different neural networks to process subliminal versus supraliminal faces has not been established, something the researchers at Osaka University aimed to address with this research.

"We are better at recognizing our own face compared to faces of others, even when the information is delivered subliminally," says lead author of the study Chisa Ota. "However, little is known about whether this advantage involves the same brain or different areas that are activated by supraliminal presentation of our face."

To address this, the researchers used [functional magnetic resonance](#) imaging (fMRI) to examine the differences between brain activity elicited by subliminally presented images of the faces of participants and faces of others. They also examined [brain activation](#) produced by subliminally presented images of faces with modified features.

"The results provided us with new insights regarding the neural mechanisms of the self-face advantage," explains Tamami Nakano, senior author. "We found that activation in the [ventral tegmental area](#), which is a central component of the dopamine reward pathway, was stronger for subliminal presentations of the participant's face compared with faces of others."

Instead, subliminal presentation of the faces of others induced activation in the amygdala of the brain, which is known to respond to unfamiliar information. This difference in [brain](#) responses to the face of the participant or those of others was consistent even when the faces were modified, as

long as the shapes of the facial features were retained.

"Our findings indicate that the dopamine reward pathway is involved in enhanced processing of one's own face even when the information is subliminal," says Tamami Nakano. "Furthermore, discrimination of one's own face from those of others appear to rely on the [information](#) of facial parts."

These findings advance the understanding of the neural mechanisms of subliminal self-facial processing. Given that the dopamine reward pathway is automatically involved in unconscious self-facial processing, this research may have applications in efforts to unconsciously manipulate motivation.

More information: Chisa Ota et al. Self-Face Activates the Dopamine Reward Pathway without Awareness, *Cerebral Cortex* (2021). DOI: [10.1093/cercor/bhab096](https://doi.org/10.1093/cercor/bhab096)

Provided by Osaka University

APA citation: Viewing your own face, even subconsciously, is rewarding (2021, April 19) retrieved 23 April 2021 from <https://medicalxpress.com/news/2021-04-viewing-subconsciously-rewarding.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.