

Once-a-week insulin treatment could be game-changing for patients with diabetes

19 April 2021



Credit: CC0 Public Domain

Treating people with Type 2 diabetes with a new once-a-week injectable insulin therapy proved to be safe and as effective as daily insulin injections, according to the results of two international clinical trials published online today in *Diabetes Care*. The studies suggest that the once-weekly treatment could provide a convenient alternative to the burden of daily insulin shots for diabetes patients.

Starting and maintaining insulin treatment remain a challenge for millions of patients worldwide with Type 2 diabetes. Fear of injections and the inconvenience and burden of injectable therapy contribute to the barriers against insulin therapy initiation and adherence. The effectiveness and safety of ongoing insulin treatment are also highly dependent on other factors, such as the accuracy of dosages, timing, and glycemic targets. Health care providers believe that reducing the frequency of treatment administration with advances, such as the once-weekly insulin used in these phase 2 trials, may decrease the reluctance to initiate insulin therapy while improving long-term adherence, glucose control, and ultimately, patient well-being.

Insulin, which has been the foundation of diabetes treatment for 100 years, is an effective glucoselowering agent and is safe when used at the correct dose," says Ildiko Lingvay, M.D., M.P.H., M.S.C.S., a professor of internal medicine and population and data sciences at UT Southwestern. "Insulin treatment is burdensome, requires frequent injections, and continues to carry a certain stigma. The development of an effective and safe insulin that can be administered once a week is a huge advance in the field."

Lingvay, who is a consultant for Novo Nordisk, is the lead author of one of the studies, which involved 205 patients from seven countries (the U.S., Croatia, Germany, Hungary, Poland, Slovakia, and Spain). The clinical trial consisted of a two-week screening period, 16 weeks of treatment, and a five-week follow-up to evaluate three different ways to adjust and optimize the insulin dose and determine which one presented the best balance between effectively lowering glucose while minimizing low-glucose events.

She also is an author of the second study that included 154 patients from five countries (the U.S., Canada, the Czech Republic, Germany, and Italy). This trial followed the same 23-week time frame and evaluated practical aspects of insulin use as well as the best ways to transition from a daily regimen to the new weekly insulin injections. The researchers determined that starting with a higher first dose—called a loading dose—allowed patients to reach their optimal glucose target faster.

"These two studies served as the steppingstones for a large phase 3 clinical trial program that is currently ongoing at UT Southwestern and other sites, which is designed to evaluate the efficacy of once-weekly insulin administration in patients with either Type 1 or Type 2 diabetes," Lingvay says. "A weekly insulin is a game-changer that will decrease the treatment burden for patients while also improving compliance. This treatment will also



decrease the burden on those who care for patients with diabetes requiring insulin. For example, for patients who need help injecting, those living in long-term care facilities, and those with memory problems, a once-weekly insulin will facilitate treatment and decrease the burden on the care providers."

More information: Harpreet S. Bajaj et al. Switching to Once-Weekly Insulin Icodec Versus Once-Daily Insulin Glargine U100 in Type 2 Diabetes Inadequately Controlled on Daily Basal Insulin: A Phase 2 Randomized Controlled Trial, *Diabetes Care* (2021). DOI: 10.2337/dc20-2877

Provided by UT Southwestern Medical Center

APA citation: Once-a-week insulin treatment could be game-changing for patients with diabetes (2021, April 19) retrieved 30 November 2022 from <u>https://medicalxpress.com/news/2021-04-once-a-week-insulin-treatment-game-changing-patients.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.