

# Restart of the Johnson & Johnson COVID-19 vaccine: A doctor explains why benefits far outweigh risks

26 April 2021, by William Petri



Credit: Unsplash/CC0 Public Domain

Editor's note: The Centers for Disease Control and Prevention and the Food and Drug Administration lifted the pause on the Johnson & Johnson COVID-19 vaccine on April 23, 2021, but the labels and [fact sheets](#) given to patients will carry a warning about the exceedingly low risk of developing blood clots. Also, close monitoring of the J&J vaccine along with the Moderna and Pfizer vaccines that were given emergency use authorization will continue. Dr. William Petri, an infectious-disease physician and immunologist at the University of Virginia School of Medicine, explains this development and why the agencies decided that the benefits of the vaccine far outweigh the risks.

## What was the concern with the Johnson & Johnson COVID-19 vaccine?

The FDA and CDC paused the use of the J&J [vaccine](#) on April 13, 2021 because of six initial cases of a rare and dangerous blood clot involving

the [cerebral venous sinuses](#), large blood vessels in the brain. This clotting, also called [thrombosis](#), can also occur in the large blood vessels of the abdomen and legs and is associated with low platelet counts, or [thrombocytopenia](#). This condition is called thrombosis with thrombocytopenia syndrome (TTS).

An unusual aspect of these clots is that they occur with a low platelet count. Platelets are cells in the blood that typically aid in clotting. TTS is likely caused by the J&J vaccination rarely causing the production of an antibody against part of a platelet called [Heparin-Platelet factor 4](#). These antibodies induce the platelets to cause clots. TTS appears similar to a rare side effect from the blood thinner [heparin](#) called [heparin-induced thrombocytopenia](#).

A panel of independent doctors and scientists known as the Advisory Committee on Immunization Practices (ACIP) extensively reviewed data after the pause. Upon further review of the total of 7 million doses of J&J vaccine administered, nine additional instances of this type of blood clot were identified, all in women.

## What did the CDC and FDA do on April 23?

The CDC and the FDA announced that it was [safe to restart the use of the Johnson & Johnson COVID-19 vaccine](#) because the benefits of protection against COVID-19 far outweigh the risks. They did so after meeting twice with the independent vaccine experts of the ACIP, which also recommended restarting.

## Why did the CDC and FDA restart the use of the Johnson & Johnson COVID-19 vaccine?

First, the side effect is rare. According to the agencies, it has occurred at most in [one in 100,000](#)

[young women](#) receiving the Johnson & Johnson vaccine. No men nor women over 50 are thought to have had this side effect. Second, the J&J vaccine is highly effective at preventing COVID-19, which is a common and severe infection that has killed 1 in 500 Americans.

Third, the J&J vaccine has advantages over the mRNA vaccines made by Pfizer and Moderna in that it requires only a single dose. It also is easier to transport and store, since it does not require super-low temperatures as do the Pfizer and Moderna vaccines.

Fourth and most importantly, now that the side effect is known, people can make informed choices about which COVID-19 vaccine to receive. A young woman on oral contraceptives, which also can promote blood clots, might logically choose one of the mRNA vaccines, whereas a man might choose the J&J vaccine since it is a single dose. Young women are the riskiest group.

### **Is this side effect seen with the other COVID-19 vaccines?**

TTS is not a side effect of the mRNA vaccines made by Moderna and Pfizer BioNTech. There have been [no cases identified in the 97.9 million doses of the Pfizer BioNTech](#) or in the 85 million doses of the Moderna vaccine. Five recipients of the mRNA vaccines had cerebral venous sinus thrombosis, but not with thrombocytopenia, so it was not TTS. TTS appears to be a side effect of only two COVID vaccines, both of which use a common cold virus called an adenovirus as part of the vaccine: the [J&J vaccine in the U.S. and the Oxford-Astrazeneca](#) vaccine used in Europe but not in the U.S.

### **How would I know if I have TTS?**

TTS has occurred only in women and in the period of six to 15 days after vaccination. Symptoms can include [severe headache, nausea, vomiting and abdominal pain followed by weakness on one side of the body](#), difficulty in speaking, seizures and loss of consciousness.

### **How is TTS treated?**

The clots from TTS are treated with what are called [directly acting anticoagulants](#), and not with heparin. This was part of the reason for the pause, so that word could get out to health care providers on how to properly treat this rare form of [blood clots](#).

### **Should I get the J&J vaccine?**

My recommendation if you are a woman under the age of 50 that you be immunized with the Pfizer or Moderna vaccines, and not the J&J vaccine. This is because this rare [side effect](#) has been observed only in [young women; 13 of the 15 women](#) who developed TTS are in this age group. For everyone else, the J&J vaccine offers the convenience of needing one and not the two doses of the other vaccines.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

APA citation: Restart of the Johnson & Johnson COVID-19 vaccine: A doctor explains why benefits far outweigh risks (2021, April 26) retrieved 9 May 2021 from <https://medicalxpress.com/news/2021-04-restart-johnson-covid-vaccine-doctor.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*