


Which medications are most toxic to the liver?

5 May 2021



drug-induced liver injury in hospitalized patients: A matched case-control study, *British Journal of Clinical Pharmacology* (2021). DOI: [10.1111/bcp.14847](https://doi.org/10.1111/bcp.14847)

Provided by Wiley

Credit: CC0 Public Domain

A new study published in the *British Journal of Clinical Pharmacology* provides insights on how common hospitalized patients develop liver injury from taking different medications.

When investigators analyzed the records of 156,570 hospitalized patients, they found 499 cases of drug-induced liver injury (DILI), for an incidence of 0.32%. Anti-infective agents, cancer medications, and [nonsteroidal anti-inflammatory drugs](#) were the major categories of drugs causing DILI, and the highest incidence was due to voriconazole (an antifungal medication). Patients with high cholesterol, [cardiovascular disease](#), pre-existing [liver disease](#), and prior surgeries faced a higher risk of DILI.

"We have carried out DILI-related real-world study with the help of information technology to provide a more accurate reference for safe and rational drug use," said corresponding author DaiHong Guo, of the Chinese People's Liberation Army General Hospital, in Beijing.

More information: Xianghao Kong et al, Incidence, characteristics and risk factors for

APA citation: Which medications are most toxic to the liver? (2021, May 5) retrieved 7 October 2022 from <https://medicalxpress.com/news/2021-05-medications-toxic-liver.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.