


Researchers find obesity linked to reduced blood flow to the brain

5 May 2021



Credit: CC0 Public Domain

A new study from scientists at The Irish Longitudinal Study on Aging (TILDA) at Trinity College Dublin reveals important findings, indicating that being overweight or obese significantly reduces blood flow in the brain. The study also shows that increased physical activity can positively modify, or even negate, this reduction in brain blood flow.

The study contains relevant information which is of great interest to the general public; since reduced blood flow in the brain, or 'cerebral hypoperfusion,' is an early mechanism in vascular dementia and Alzheimer's disease.

Obesity and health challenges

According to the World Health Organization (WHO), obesity is a worsening [health](#) crisis that has reached epidemic proportions globally, with over 1 billion adults overweight—and at least 300 million clinically obese. It continues to be a major contributor to global rates of chronic disease and disability, affecting overall quality of life, while placing increased strain on the immune system

which is of the utmost importance given the current COVID-19 situation. Obesity is also a significant public health concern given its [negative impact](#) on physiological function, especially as we age. Finding easily implemented and cost-effective ways to tackle the impact of obesity is particularly important to help protect against negative health outcomes in later life.

What are the findings of the TILDA study?

The study investigates three different measures of obesity—body mass index (BMI), waist-to-hip ratio and waist circumference, as well as [physical activity](#), in adults over 50 years.

Brain blood flow was measured using cutting-edge MRI scanning and analysis techniques. The findings reveal that being overweight or obese is associated with reduced blood supply to the brain. Whereas [brain blood flow](#) is known to decline with age, in this study the negative influence of obesity on brain blood flow was shown to be greater than that of age. However, being physically active helps to cancel out the negative effects of obesity on brain blood flow.

Key Findings

- Increased BMI, waist-to-hip ratio, and waist size are associated with less blood supply to the brain.
- A waist size increase of +1cm is associated with the same reduction in brain blood flow as +1 year of age.
- Higher levels of physical activity modify the associations between reduced brain blood flow and obesity.

Recommendations

The study recommends at least 1.5 to two hours of 'being active' throughout the day, engaging in activities that require moderate effort. These

include activities that cause one to breathe harder than normal, such as fast walking or cycling. However, any increase in physical activity, particularly if integrated into daily or weekly habits, such as gardening, should help maintain and potentially improve brain blood flow.

Dr. Silvin Knight, Research Fellow at TILDA and lead author, said:

Consistent, healthy blood supply to the brain is critical, as it ensures that the brain is provided with enough oxygen and nutrients to function correctly. If brain blood flow becomes impaired, it can lead to serious health issues as we age, such as increasing the risk of dementia and Alzheimer's disease. We know that obesity can predispose a person to age-related conditions, illness, and disease, and even reduce life expectancy by up to six years in men and seven years in women, after the age of forty. Our study reveals clear associations between obesity and reduced [blood supply](#) to the brain in an older population. The study also shows the importance of being physically active for older overweight or obese individuals, as this may help to protect against reduced brain blood flow and the poor health outcomes that can arise from this.

Professor Rose Anne Kenny, Principal Investigator of TILDA, and co-author of the study, said: "Many experts have shown that obesity and aging have very similar effects on the biology of aging; diseases associated with obesity are similar to those of aging and age-related diseases—heart disease, diabetes, high [blood](#) pressure, kidney failure, arthritis, susceptibility to infections—including COVID-19."

Our study not only shows that there is a link between obesity and reduced [brain blood flow](#), but also that it is possible to protect against the negative consequences of obesity through regular physical exercise. Whereas these findings are of relevance in the global context, because of the rapidly evolving global burden of obesity, the research is especially important to Irish adults because obesity and being overweight is a considerable health issue in Ireland.

Previous TILDA research has shown that over one-third of Irish adults aged 50 and older are obese and a further 43% overweight. As we prepare our society for a growing aging population, we can use this evidence to prepare meaningful public health policies that will promote impactful and positive lifestyle habits, such as regular physical activity, to mitigate against some of the negative consequences of the growing [obesity](#) crisis.

More information: Silvin P. Knight et al. Obesity is associated with reduced cerebral blood flow – modified by physical activity, *Neurobiology of Aging* (2021). [DOI: 10.1016/j.neurobiolaging.2021.04.008](https://doi.org/10.1016/j.neurobiolaging.2021.04.008)

Provided by Trinity College Dublin

APA citation: Researchers find obesity linked to reduced blood flow to the brain (2021, May 5) retrieved 2 December 2022 from <https://medicalxpress.com/news/2021-05-obesity-linked-blood-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.