

Older adults having surgery less frequently—but it depends on the surgery and hospital

10 May 2021



Credit: Pixabay/CC0 Public Domain

Many have assumed that the rates of major abdominal surgeries in adults over 65 is increasing over time as the U.S. population ages and as new technology renders surgical procedures safer for older adults. Contrary to this popular belief, a new study from the University of Chicago Medicine found the frequency of abdominal surgery in older adults is decreasing, especially among adults over the age of 85. The study, which examined data from 2002 to 2014, was published May 10 in the *Journal of the American Geriatric Society*.

While the research was not able to determine the exact reasons for this shift, the results indicate that improvements in [medical treatments](#) and cancer screening for older adults may be reducing the need for invasive surgeries, which can have negative impacts on patient's long-term quality of life.

"There was a really stark decrease in the number of surgeries among older adults over the age of 65

throughout the study period," said first author Daniel Rubin, MD, Associate Professor of Anesthesia and Critical Care at UChicago Medicine. "This suggests we're getting better at determining who would benefit most from a [surgery](#), and also possibly that we've developed better and less invasive alternative treatments."

Rubin was inspired to investigate this question after reading a 2019 New York Times article discussing "routine" surgeries in adults over 85.

"This is a very personal topic for me because I was very close to my grandparents, and I saw their experiences with geriatric surgeries," said Rubin. "My grandmother had an emergency hysterectomy at 92. Despite a stay in the intensive care unit and delirium she was able to go home and live independently afterward. My 89-year-old grandfather, on the other hand, declined a Whipple procedure for treating pancreatic cancer. So I could see early on how these big surgical interventions can really affect older adults."

While the researchers found the overall frequency of surgery in older adults is decreasing, they did note some exceptions.

"The biggest surprise was that we saw a redistribution of surgical procedures to certain types of hospitals," Rubin said. "There was a large shift away from rural and non-teaching urban hospitals to major academic medical centers; these were the only hospitals that saw an increase in the frequency of procedures performed on older adults. We think this may be due to increased specialization and that some procedures that may not be available at a smaller or rural institution are more likely to be offered at a larger, academic hospital."

Another trend that was less surprising was the

increase in certain procedures, such as the Whipple procedure for treating pancreatic cancer. "With [pancreatic cancer](#), there aren't really many less-invasive options available, and outcomes are generally good even for older adults," Rubin said. "We perform Whipple procedures on 85- or 90-year-old adults here at UChicago Medicine simply because it is usually the most effective option."

Provided by University of Chicago Medical Center

The researchers say these results are important because older adults often struggle to recover after major surgeries. The study found that over half of adults over 85 will require at least some transitional, post-acute medical care after a surgical procedure.

"Older adults really want to get back to their normal life after an operation," said Rubin. "But recovery can be difficult, and the rates at which older adults require post-acute care—such as time in a rehabilitation facility or long-term care—is high. These results can help inform the decisions made by patients and providers on whether or not a surgery is the right choice for them."

Understanding how trends in surgical procedures change over time can help illuminate which options are the most likely to lead to optimal outcomes—and determine which patients will most benefit from specific interventions.

"We're very good at understanding overall mortality for certain procedures, but not as good at identifying which individual patients will have poor outcomes," said Rubin. "It's important for patients and their families to understand that these surgeries can lead to a challenging recovery trajectory for older adults, even if the surgery goes well. My hope is that this research, and other studies like it, will help us better understand how to help [older adults](#) prepare for major surgeries, and help us identify which individual patients will benefit the most from surgical intervention."

More information: Daniel S. Rubin et al, U.S. trends in elective and emergent major abdominal surgical procedures from 2002 to 2014 in older adults, *Journal of the American Geriatrics Society* (2021). [DOI: 10.1111/jgs.17189](https://doi.org/10.1111/jgs.17189)

APA citation: Older adults having surgery less frequently—but it depends on the surgery and hospital (2021, May 10) retrieved 1 June 2021 from <https://medicalxpress.com/news/2021-05-older-adults-surgery-frequentlybut-hospital.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.