


Rare COVID-19 response in children explained

17 May 2021, by Bess Connolly



Credit: Unsplash/CC0 Public Domain

One of the enduring mysteries of the COVID-19 pandemic is why most children tend to experience fewer symptoms than adults after infection with the coronavirus. The immune system response that occurs in the rare cases in which children experience life-threatening reactions after infection may offer an important insight, a Yale-led study published in the journal *Immunity* suggests.

While many children infected with the [virus](#) are asymptomatic or go undiagnosed, about one in 1,000 children experience multi-system [inflammatory response](#) (MIS-C) four to six weeks after confirmed infection with SARS-CoV-2, the virus that causes COVID-19. The condition is marked by a variety of symptoms, including fever, abdominal pain with vomiting and/or diarrhea, rash, and cardiovascular and neurological problems. If diagnosed early, the condition is readily treatable with immune suppressants such as steroids. If left untreated, however, it can be fatal.

"Why does this happen when there is no virus or anti-viral response still present and in kids? And why is it only occurring in youth?" asked Carrie Lucas, an assistant professor of immunobiology at

Yale and corresponding author of the new study.

In an exhaustive analysis, Lucas and her lab tested blood from children with MIS-C, adults with severe COVID symptoms, as well as healthy children and adults. They found that children with MIS-C had immune system signatures distinct from other groups.

Specifically, the children with MIS-C had high levels of alarmins, molecules that make up part of the innate immune system which is mobilized quickly to respond to all infections. Other research findings have suggested that a child's innate [immune system response](#) may be stronger than those of adults, one possible explanation for why they generally experience milder symptoms than adults after [infection](#).

"Innate immunity may be more active in children who are infected with virus," Lucas said. "But on the flip side, in rare cases it may get too revved up and contribute to this inflammatory disease."

Children diagnosed with MIS-C were also found to have a marked elevation of certain adaptive immune responses, which are defenses to combat specific pathogens—such as the virus causing COVID-19—and that typically confer immunological memory. But instead of being protective, the responses produced in these affected children appear to attack a variety of host tissues, a hallmark of autoimmune diseases.

Lucas speculates that the initial immune response in these rare cases triggers a cascade that damages healthy tissue, which in turns makes the tissue more susceptible to attack by autoantibodies.

In the meantime, the peculiar immune system signatures of MIS-C could help in the diagnosis and early treatment options of [children](#) at high risk of the disorder, Lucas said.

More information: Anjali Ramaswamy et al.
Immune dysregulation and autoreactivity correlate
with disease severity in SARS-CoV-2-associated
multisystem inflammatory syndrome in children,
Immunity (2021). DOI:
[10.1016/j.immuni.2021.04.003](https://doi.org/10.1016/j.immuni.2021.04.003)

Provided by Yale University

APA citation: Rare COVID-19 response in children explained (2021, May 17) retrieved 2 September 2022 from <https://medicalxpress.com/news/2021-05-rare-covid-response-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.