

People of color twice as likely to die after traumatic brain injury, new study finds

3 June 2021, by Erik Robinson



Credit: Unsplash/CC0 Public Domain

People of color are more than twice as likely to die after a traumatic brain injury as white people, according to a new retrospective review from Oregon Health & Science University.

The study published today in the journal *Frontiers in Surgery*.

In the report, "Racial and Ethnic Inequities in Mortality During Hospitalization for Traumatic Brain Injury: A Call to Action," the researchers analyzed more than a decade of data related to the [health outcomes](#) and demographics of thousands of patients treated for traumatic head injuries at OHSU Hospital, one of two Level 1 trauma centers in the state.

They found a clear delineation of worse outcomes for people of color.

"We have a societal and professional duty to recognize and accept that the effects of structural racism have taken hold of our patients' [health](#) long before they arrive in our trauma bays, ICU beds, and operating tables," they write. "These disparities permeate our society and contribute to

inequitable health outcomes, and we must take action to identify the factors which perpetuate this disproportionate suffering."

The researchers found no bias in the treatment patients received while in the hospital. Measurements including the number of days spent in [intensive care](#) and charges for hospital services were similar between groups.

Rather, they say the findings highlight underlying disparities in health that disproportionately affect people of color.

"If you ran the same analysis with patients with pneumonia, you might find the same results," said senior author Ahmed M. Raslan, M.D., FAANS, associate professor of neurological surgery in the OHSU School of Medicine. "Our findings point to the problem in a more robust and clear way. The problem is rooted in social determinants of health, including factors such as diet, lifestyle, occupations and access to [health care](#)."

The study is the first to review racial and ethnic disparities in the field of neurosurgical trauma.

Researchers reviewed outcomes for 6,352 patients treated for traumatic brain injuries between 2006 and 2017. The overall mortality rate was 9.9% among all patients. Among the 1,504 patients who identified as racial or ethnic minorities, 14.6% died.

Weighted for severity of injury, age and other factors, researchers calculated an in-hospital mortality hazard ratio of 2.21 for minority patients compared to white patients.

"Dr. Raslan's research is incredibly important in highlighting disparities in health outcomes experienced by people of color," said Nathan Selden, M.D., Ph.D., professor and chair of neurological surgery in the OHSU School of Medicine. "Together with our department's efforts to

improve the care of traumatic brain injuries, and more broadly to improve diversity in medicine, I am confident OHSU can really make a difference."

More information: Emma A. Richie et al, Racial and Ethnic Inequities in Mortality During Hospitalization for Traumatic Brain Injury: A Call to Action, *Frontiers in Surgery* (2021). DOI: [10.3389/fsurg.2021.690971](https://doi.org/10.3389/fsurg.2021.690971)

Provided by Oregon Health & Science University

APA citation: People of color twice as likely to die after traumatic brain injury, new study finds (2021, June 3) retrieved 4 November 2022 from <https://medicalxpress.com/news/2021-06-people-die-traumatic-brain-injury.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.