

# New drug could help halt dementia after multiple head injuries

June 7 2021

---



Credit: Unsplash/CC0 Public Domain

A world-first international study led by the University of South Australia has identified a new drug to stop athletes developing dementia after sustaining repeated head injuries in their career.

The link between concussion and neurodegenerative diseases is well established, but new research findings could halt the progression of [chronic traumatic encephalopathy](#) (CTE) in sportspeople who sustain repeated blows to the head.

CTE is a progressive and [fatal brain disease](#) associated with the accumulation of a protein known as hyperphosphorylated tau which affects cognition and behavior.

In a paper published in *Scientific Reports*, UniSA Emeritus Professor Bob Vink and colleagues show how repeated concussions can cause CTE and a way to block it with a specially developed drug.

The findings will potentially have significant implications for athletes who play contact sports—such as boxers and footballers—as well as military veterans sustaining [head injuries](#) in conflict.

The team of researchers from Adelaide, Melbourne and the United States say the brain releases a neurotransmitter called substance P in the event of a head [injury](#), causing abnormal amounts of the tau protein to collect inside neurons.

"Tau protein tangles are a feature of CTE, which reportedly leads to memory problems, confusion, personality changes, aggression, depression and suicidal thinking," Prof Vink says.

"Our research shows that by blocking substance P with a specific drug, we can prevent the tau protein tangles from developing in the brain and causing neurological problems."

The treatment was successfully tested in animal models, giving hope that CTE can be prevented in humans.

Prof Vink says the next step is [human clinical trials](#), but that could take several years given that currently CTE can only be diagnosed post-mortem.

A study of 14,000 Americans over 25 years, published in *Alzheimer's and Dementia* in March, showed that people who sustained even one [head](#) injury were 25 percent more likely to develop dementia later in life. This risk increased with multiple traumatic brain injuries.

[The Guardian](#) also reported in April that an analysis of late AFLW player Jacinta Barclay's brain uncovered neurological damage at age 29, highlighting the risks of repeated concussions to both sexes. Previous research has focused on the impact of [brain](#) injuries in male athletes, but females are more likely to sustain concussions.

**More information:** Frances Corrigan et al, NK1 antagonists attenuate tau phosphorylation after blast and repeated concussive injury, *Scientific Reports* (2021). [DOI: 10.1038/s41598-021-88237-0](https://doi.org/10.1038/s41598-021-88237-0)

Andrea L. C. Schneider et al, Head injury and 25-year risk of dementia, *Alzheimer's & Dementia* (2021). [DOI: 10.1002/alz.12315](https://doi.org/10.1002/alz.12315)

Provided by University of South Australia

Citation: New drug could help halt dementia after multiple head injuries (2021, June 7) retrieved 24 November 2023 from <https://medicalxpress.com/news/2021-06-drug-halt-dementia-multiple-injuries.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--