

New target identified for rapid antidepressant drugs

November 10 2021, by Aaron Conley



Credit: Unsplash/CC0 Public Domain

Vanderbilt researchers found that ketamine's rapid antidepressant action is due to specific synaptic effects. This represents a new target for drug development that could fill a major gap in care for depression.

The research was led by Lisa Monteggia, professor of pharmacology and director of the Vanderbilt Brain Institute, Ege Kavalali, professor and acting chair of the Department of Pharmacology and William Stokes Professor of Experimental Therapeutics, and first author Kanzo Suzuki, [research](#) instructor in pharmacology.

"We examined the functional role of eEF2K, a protein involved in ketamine's rapid antidepressant action, in neurons," Monteggia said. "We found that inducing similar synaptic effects independent of eEF2K resulted in rapid behavioral effects analogous to ketamine in preclinical models. This is the first study that directly examines whether the synaptic effects are producing the antidepressant behavioral activity of ketamine."

Uncovering this pathway is a significant step toward understanding ketamine's action in the brain and exploring whether a new class of therapeutics targeting this pathway can provide rapid antidepressant treatment.

Why it matters

Depression, which the World Health Organization says costs \$1 trillion annually in lost productivity, affects 264 million people around the world. Current pharmacological treatments for depression typically take several weeks to show efficacy and are not effective in approximately half of patients. Individuals whose depression is treatment-resistant are at far higher risk of suicide.

Monteggia and Kavalali's goal is to find therapeutics that can overcome treatment-resistant depression and reduce [depression](#)-related loss of life.

What's next?

Kavalali, Monteggia and their labs plan to further probe these specific synaptic effects required for rapid antidepressant activity. In particular, they aim to link the synaptic changes to behavioral phenotypes by systematically tracing the impact of these changes on neuronal circuits.

More information: Kanzo Suzuki et al, Convergence of distinct signaling pathways on synaptic scaling to trigger rapid antidepressant action, *Cell Reports* (2021). [DOI: 10.1016/j.celrep.2021.109918](https://doi.org/10.1016/j.celrep.2021.109918)

Provided by Vanderbilt University

Citation: New target identified for rapid antidepressant drugs (2021, November 10) retrieved 5 October 2023 from <https://medicalxpress.com/news/2021-11-rapid-antidepressant-drugs.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--