

Disinfectant use during pregnancy linked to childhood asthma and eczema

March 28 2022



Credit: Pixabay/CC0 Public Domain

Use of disinfectants by pregnant women may be a risk factor for asthma and eczema in their children, finds a population study, published online in *Occupational & Environmental Medicine*.

Disinfectants are used frequently in hospitals and other [medical facilities](#), and the COVID-19 pandemic has led to an increase in their use in medical settings and also more widely, including by the general population.

Exposure to [disinfectants](#) in the workplace has been linked to asthma and dermatitis previously in the workers exposed, but few studies have looked at the impact of disinfectant use during pregnancy and the subsequent development of allergic disease in children.

The authors used data on 78 915 [mother-child pairs](#) who participated in the Japan Environment and Children's Study to examine whether mothers' exposure to disinfectants in the workplace was associated with increased risk of diagnosis of allergic diseases in their children when aged 3 years.

The odds of children having asthma or eczema were significantly higher if their mothers used disinfectant one to six times a week compared with the odds in children of mothers who never used disinfectants.

There was an exposure-dependent relationship between [prenatal exposure](#) to disinfectants and the odds of children experiencing these

allergic conditions, with the children of mothers exposed to disinfectants every day having the highest odds of a diagnosis—26% greater for asthma and 29% greater for eczema than children of mothers who were never exposed to disinfectants.

There were no significant associations between disinfectant use and food allergies.

This is an observational study, and as such, can't establish cause. The authors also noted some limitations, including that the information on disinfectant use by mothers was self-reported with specific disinfectants not identified. Diagnoses of allergic diseases in children were also reported by mothers.

Nevertheless, the authors conclude, "Our findings indicate that exposure [to disinfectants] during pregnancy exerts an effect on allergies in offspring regardless of whether the mother returns to work when the child is 1 year old, and suggest an effect by exposure during pregnancy alone."

They add, "Given the current increased use of disinfectants to prevent new coronavirus infections, it is of great public health importance to consider whether prenatal disinfectant exposure is a risk for the development of allergic diseases."

Several mechanisms that could explain the increased risk of allergic disease in children following their mothers' exposure to disinfectants during pregnancy were suggested by the authors.

They included microbiome-mediated (disinfectants impact the gut and skin microflora of the mother and subsequently the child), immune-mediated (exposure to some [chemical compounds](#) during pregnancy impacts the [immune response](#) in the fetus), postnatal exposure ([children](#)

inhaled or touched molecules of disinfectant on the skin of their mothers), or bias (mothers who use medical disinfectants frequently are likely to be more medically knowledgeable and have better access to healthcare).

More information: Prenatal occupational disinfectant exposure and childhood allergies: the Japan Environment and Children's study, *Occupational & Environmental Medicine*, 2022. [DOI: 10.1136/oemed-2021-108034](https://doi.org/10.1136/oemed-2021-108034)

Provided by British Medical Journal

Citation: Disinfectant use during pregnancy linked to childhood asthma and eczema (2022, March 28) retrieved 11 July 2023 from <https://medicalxpress.com/news/2022-03-disinfectant-pregnancy-linked-childhood-asthma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.