

Inhaled antibiotics could help reduce antimicrobial resistance

May 13 2022



Credit: Pixabay/CC0 Public Domain

Using inhaled antibiotics to treat lower respiratory tract infections could help reduce antimicrobial resistance, according to researchers from the

Universities of Bristol and Bath.

Acute [lower respiratory tract infections](#) (aLRTIs) are the most common condition managed in primary care worldwide and antibiotic resistant strains are the leading cause of antimicrobial deaths. Oral antibiotics are currently inappropriately given in around 50% of cases, contributing to antimicrobial resistance and medicines waste.

In an article published in *The Lancet Respiratory Medicine* (11 May), the researchers propose that inhaled antibiotics could be effective in treating people seen in [primary care](#), particularly those with long term lung conditions such as asthma and COPD ([chronic obstructive pulmonary disease](#)). These patients are already familiar with using inhalers, and inhaled antibiotics could deliver higher concentrations in smaller doses than when taken by mouth, and therefore cause less harm.

Professor Alastair Hay from the Centre for Academic Primary Care, University of Bristol and lead author of the article, said: "The evidence base for using of inhaled antibiotics is really lacking, but the evidence that is available suggests that they could be a game-changer when it comes to the treatment of acute lower respiratory tract infections. It's vital that we consider all possible avenues to reduce antimicrobial resistance, which remains a major threat to public health globally.

"A lot of damage can be done through the inappropriate use of oral antibiotics, including side effects such as diarrhea, vomiting and skin rashes, and damage to healthy gut bacteria that are indiscriminately destroyed, allowing resistant bacteria to flourish. Inhaled [antibiotics](#) could reduce side effects and these so-called 'bystander' [antimicrobial resistance](#) effects, by ensuring only the affected part of the body is treated.

"We are calling on governments around the world to make funds

available for companies to develop new drug delivery systems so that patients can reduce their use of [oral antibiotics](#) and thereby reduce the risks of developing side effects and acquiring antibiotic resistant bacteria."

More information: Alastair D Hay et al, Inhaled antibiotics for acute lower respiratory tract infections in primary care: a hypothesis, *The Lancet Respiratory Medicine* (2022). [DOI: 10.1016/S2213-2600\(22\)00172-2](#)

Provided by University of Bristol

Citation: Inhaled antibiotics could help reduce antimicrobial resistance (2022, May 13) retrieved 2 March 2023 from <https://medicalxpress.com/news/2022-05-inhaled-antibiotics-antimicrobial-resistance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.