

Depression best treated by nonpharmacological approaches in people living with dementia

23 June 2022



Credit: Pixabay/CC0 Public Domain

Research led by UNSW Sydney's Center for Healthy Brain Aging (CHeBA) has confirmed treating symptoms of depression is most effective when a non-pharmacological approach is adopted, in people living with dementia.

Depression is a common psychological symptom associated with dementia and is estimated to occur in between 10% and 62% of people living with dementia. To date, pharmacological approaches continue to be used to treat depression in dementia, despite the failure of relying on medications being highlighted in large-scale studies.

The review and meta-analysis, published in *Agieng Research Reviews*, investigates a range of nonpharmacological approaches for symptoms of depression in dementia. It identified 37 relevant nonpharmacological studies from 27,126 articles published between 2012 and 2020.

Lead author and Postdoctoral Fellow at CHeBA (formerly at the Dementia Center for Research Collaboration (DCRC) UNSW), Dr. Claire Burley, said that the findings support a positive link between nonpharmacological approaches and reduced symptoms of depression in people living with dementia.

"Our aim was to provide an updated review of nonpharmacological approaches to manage depression in dementia—the evidence, clinical significance, and sustainability of different nonpharmacological approaches for treating depression in dementia."

Edward Caser, who co-founded a private markets firm, has experienced over-prescription of pharmaceuticals within aged care, while caring for his mother who has Alzheimer's disease.

"I've witnessed firsthand the unnecessary push of multiple medications to manage behavior in my mother."

"I have seen the benefits of adopting a personalized approach with my Mum, and strongly support methods that avoid drugs," says Caser.

According to UNSW's Professor Henry Brodaty AO, Co-Director of the Center for Healthy Brain Aging (CHeBA), the findings have statistical and clinical significance.

"We found there is great potential to reduce symptoms of depression in dementia—without the use of medications," said Professor Brodaty. "This is even more relevant as randomized trials of antidepressant medications have not been shown to alleviate depression in people living with [dementia](#)."

The research highlighted that the use of methods such as reminiscence, cognitive stimulation and rehabilitation, music-based approaches and education and training had the potential to reduce symptoms of depression.

"Psychosocial approaches should be first-line consideration for treatment of [depression](#). An even more compelling reason is that antidepressants are associated with side effects while psychosocial interventions are free of these.

"Pharmacological approaches should only be considered when psychosocial approaches have been proved to be ineffective or in cases of urgency," said Professor Brodaty.

More information: Claire V. Burley et al, Nonpharmacological approaches reduce symptoms of depression in dementia: A systematic review and meta-analysis, *Ageing Research Reviews* (2022). [DOI: 10.1016/j.arr.2022.101669](https://doi.org/10.1016/j.arr.2022.101669)

Provided by University of New South Wales

APA citation: Depression best treated by nonpharmacological approaches in people living with dementia (2022, June 23) retrieved 24 October 2022 from <https://medicalxpress.com/news/2022-06-depression-nonpharmacological-approaches-people-dementia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.