

# Gender differences in behaviors linked to faster cognitive decline

June 30 2022

---



Credit: Pixabay/CC0 Public Domain

Men who experience behavior changes including apathy or having false beliefs and perceptions in later life are at risk of faster cognitive decline than women, according to new research.

A study led by Charles University, in collaboration with the University of Exeter and King's College London, looked at changes in behavior in cognitively healthy people aged 50 and older, which have previously been linked to a higher risk of developing brain problems.

These behavioral changes are collectively known as mild behavioral impairment (MBI), and encompass symptoms across five domains: apathy or social withdrawal; changes in mood and anxiety; being more impulsive than usual; being socially inappropriate; and having false beliefs or hallucinations. Observing these changes may help physicians and researchers to identify people at a higher risk of developing dementia later in life.

Published in the *Journal of Alzheimer's Disease*, the study is the first to investigate gender differences in the link between these behavioral changes and brain decline. The researchers found that men who exhibit MBI symptoms are at risk of faster [cognitive decline](#) compared to women, particularly when [psychotic symptoms](#) or symptoms of decreased motivation occur.

Dr. Katrin Wolfová, Ph.D. student at Charles University in Prague, is first author of the study. She said: "Observing change in certain behavioral traits can help us to identify people who might already have a neurodegenerative disease before their cognitive functions decline. Being aware of differences between men and women in risk of developing dementia based on these behaviors is important for clinicians as well as researchers."

Previous research suggests that possible explanations of these findings might be related to structural differences in entorhinal and hippocampal regions of brain, functional differences in serotonergic system, or unequal distribution of types of dementia in men and women.

The research team, which also involved collaborators from Canada, studied more than 8,000 people enrolled in the British online cohort ([PROTECT](#)) study, led by the University of Exeter and King's College London. The participants were on average 63 years old and 73% of them were women.

To assess cognitive functions, participants completed a series of online tests. Behavioral changes, or MBI, were also assessed online via a specialized questionnaire in the PROTECT platform.

Dr. Byron Creese, of the University of Exeter, said: "This research adds to the growing evidence that some people who experience suspicious thoughts or become withdrawn in later life have more memory and thinking problems. The findings are important because they suggest that monitoring these symptoms could help identify people with memory problems early on. The research also highlights the importance of [gender differences](#) in these relationships."

Dr. Wolfová added that although there is currently no cure, patients might benefit from [early diagnosis](#) through secondary prevention and development of a comprehensive care plan. Identification of patients with preclinical Alzheimer's disease is extremely important for [clinical trials](#), as researchers need to screen potential participants with expensive biomarker tests before they can enroll them in a study. Screening with MBI to assess whether candidates are suitable for a trial might save a lot of money, and optimization of this process to reflect differences between men and women could further increase precision and reduce costs.

**More information:** Katrin Wolfova et al, Gender/Sex Differences in the Association of Mild Behavioral Impairment with Cognitive Aging, *Journal of Alzheimer's Disease* (2022). [DOI: 10.3233/JAD-220040](https://doi.org/10.3233/JAD-220040)

Provided by University of Exeter

Citation: Gender differences in behaviors linked to faster cognitive decline (2022, June 30)  
retrieved 28 February 2023 from <https://medicalxpress.com/news/2022-06-gender-differences-behaviors-linked-faster.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.