


When friendly forces become foes: Scientists blunt the impact of natural killer cells to improve vaccine effectiveness

25 July 2022, by Delthia Ricks



PD-1 dependent NK cell regulation of CD8⁺T-cells. C57BL/6 mice were infected with an adenoviral vector encoding the HBV genome and treated with anti-NK1.1 (αNK) or isotype control antibody prior to therapeutic vaccination. Intrahepatic lymphocytes were harvested 14 days after immunization. (A) Quantitative real-time PCR analysis of HBsAg mRNA extracted from the liver of infected mice. (B) Representative plot of CD8⁺T-cells isolated from the spleen of CD45.2 PD-1KO or CD45.1 wild-type (WT) mice transferred to opposite congenic mouse recipient one day prior to therapeutic vaccination. Example of PD-1 expression and IFN γ production on transferred PD-1KO and WT CD8⁺ T-cells. *, p

APA citation: When friendly forces become foes: Scientists blunt the impact of natural killer cells to improve vaccine effectiveness (2022, July 25) retrieved 8 September 2022 from <https://medicalxpress.com/news/2022-07-friendly-foes-scientists-blunt-impact.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.