

New prognostic marker discovered for multiple sclerosis severity

August 10 2022



Credit: Unsplash/CC0 Public Domain

It is essential to assess the severity of multiple sclerosis (MS) in order to choose appropriate therapeutic measures, but this cannot be reliably done using existing methods. A MedUni Vienna study now shows for the

first time that the retina can be used as a prognostic marker. Analyses revealed that retinal layer thinning as a result of an MS relapse predicts the severity of future relapses and, hence, the likelihood of disability. The results of the study have now been published in *Neurology*.

Researchers led by Gabriel Bsteh and Thomas Berger from the Department of Neurology at MedUni Vienna/University Hospital Vienna, working in collaboration with the Department of Ophthalmology and Optometrics at MedUni Vienna/University Hospital Vienna, studied 167 MS patients over a period of more than three years. They hypothesized that [retinal damage](#) due to relapse reflects the extent of damage in the brain. As the scientific analyses confirmed, the loss of approximately 5 µm (micrometers) of retinal layer thickness after [optic neuritis](#) equates to a doubling of the risk of permanent disability after the next relapse.

These predictions could be used as the basis for [treatment decisions](#) in future: the results of the study suggest that more aggressive treatment is indicated where there is significant retinal layer thinning than would be the case for a smaller degree of thinning. This is true even if the patient has no disability or only slight disability at the time of measurement.

Prognostic technique already available

The researchers used [optical coherence tomography](#) (OCT) to measure retinal layer thickness. OCT is an [imaging technique](#) that uses [infrared light](#) to produce high-resolution three-dimensional images of very thin layers of tissue in the micrometer range (1 micrometer=1 thousandth of a millimeter). It is already used as a tool for diagnosing eye diseases such as glaucoma, and for evaluating [disease progression](#).

"The technique for predicting the course of MS is therefore already available to us," said Gabriel Bsteh, first author of the study. "As we

discovered in the course of our clinical trial, measurements should be taken at initial diagnosis, directly when optic neuritis occurs in relapsing MS, and six months thereafter."

The retina as a window to the brain

Multiple sclerosis is a chronic inflammatory autoimmune disease that leads to the loss of axons and neurons throughout the entire nervous system. Although this damage often goes unnoticed by patients at first, its extent determines the prognosis for the severity of the disease. Since predictions about the course of the disease are important in MS for selecting the appropriate treatment, [medical research](#) has long been searching for reliable prognostic tools.

"In retinal layer thickness, we have found a new biomarker that represents a window to the brain, as it were," said Gabriel Bsteh, summarizing the essence of the study. If the results are confirmed in larger follow-up studies, the technique could also be applied in routine clinical practice.

More information: Gabriel Bsteh et al, Retinal Layer Thinning After Optic Neuritis is Associated With Future Relapse Remission in Relapsing Multiple Sclerosis, *Neurology* (2022). [DOI: 10.1212/WNL.0000000000200970](#)

Provided by Medical University of Vienna

Citation: New prognostic marker discovered for multiple sclerosis severity (2022, August 10) retrieved 8 October 2023 from <https://medicalxpress.com/news/2022-08-prognostic-marker-multiple-sclerosis-severity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.