

# Vitamin D supplementation may aid new-onset pediatric type 1 diabetes

26 August 2022



A1c at a mean rate of change of 0.30 percent every three months versus 0.77 percent every three months for the [placebo group](#).

"We recommend a baseline estimation of 25(OH)D concentration at the time of diagnosis of type 1 [diabetes](#), and to begin vitamin D supplementation if serum 25(OH)D concentration is

High-dose vitamin D supplementation in pediatric patients with new-onset type 1 diabetes may reduce complications, according to a study published online Aug. 18 in *Frontiers in Endocrinology*.

Benjamin Udoka Nwosu, M.D., from the Zucker School of Medicine at Hofstra/Northwell in New Hyde Park, New York, randomly assigned 36 children and adolescents with type 1 diabetes to receive either vitamin D2 (ergocalciferol, given as 50,000 international units per week for two months and then every other week for 10 months) or a placebo.

The researchers found that vitamin D was significantly associated with a lower temporal rise in hemoglobin A1c at a mean rate of change of 0.14 percent every three months versus 0.46 percent every three months for the placebo group. Additionally, vitamin D was significantly associated with the functional marker of partial clinical remission, the insulin-dose adjusted hemoglobin

APA citation: Vitamin D supplementation may aid new-onset pediatric type 1 diabetes (2022, August 26) retrieved 30 August 2022 from <https://medicalxpress.com/news/2022-08-vitamin-d-supplementation-aid-new-onset.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*