

Heavy smoking and smoke inhalation may worsen impact of heart attacks

August 26 2022



Credit: Unsplash/CC0 Public Domain

The risk of death or poor prognosis after a heart attack is more than 20-fold higher in smokers with exhaled carbon monoxide levels above 13 ppm, indicating heavy smoking and inhalation of smoke. That's the

finding of late-breaking research presented at ESC Congress 2022.

"The amount of carbon monoxide in the breath is directly linked to the number of cigarettes you smoke but also how you smoke," said principal investigator Professor Patrick Henry of Hospital Lariboisiere, Paris, France. "Individuals can inhale smoke deeply or not at all, and they may either smoke cigarettes from beginning to end or let the cigarettes burn in the ashtray. Finally, the location in which [smoking](#) occurs may play a crucial role; smoking in a small, non-ventilated room likely increases carbon monoxide concentration compared to smoking outside."

The combustion of cigarettes produces carbon monoxide, a highly toxic gas also found in motor vehicle exhaust, pollution, and malfunctioning furnaces. Carbon monoxide takes the place of oxygen in the blood and can be lethal. Professor Henry says that "patients with acute cardiac events such as heart attacks have insufficient oxygen in their coronary arteries. We hypothesized that if part of the oxygen was replaced by carbon monoxide, the event could be more severe."

During a two week period in April 2021, expiratory carbon monoxide was measured within two hours of admission in all consecutive adults hospitalized for acute cardiac events in 39 intensive cardiac care units (ICCU) in France. A total of 1,379 patients were studied. The average age was 63 years and 70% were men. Regarding the reason for admission, 720 (52%) patients had [acute coronary syndrome](#), 186 (13%) had [acute heart failure](#), and 473 (34%) had other acute cardiac conditions. The median stay in the ICCU was five days.

Patients were asked about smoking status. One-third of participants (33%) were non-smokers, 39% were former smokers, and 27% were active smokers. Carbon monoxide level was similar in non-smokers and former smokers (mean 3.6 and 3.3 ppm, respectively; $p=0.12$) and significantly higher in active smokers (mean 9.9 ppm; p

Citation: Heavy smoking and smoke inhalation may worsen impact of heart attacks (2022, August 26) retrieved 15 July 2023 from <https://medicalxpress.com/news/2022-08-heavy-inhalation-worsen-impact-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.