

Cardiovascular disease risk in adults with compromised bone health

September 7 2022



Credit: Unsplash/CC0 Public Domain

These risk factors included male sex, older age, smoking, alcohol consumption, atrial fibrillation, use of anti-hypertensive medications, prior heart attack or stroke, established cardiovascular disease, low kidney function, high systolic blood pressure, elevated cholesterol level,

and use of multiple concomitant medicines.

"Although there are some calculators to produce risk estimates of cardiovascular disease, these are not targeted at those at high risk of fracture," said corresponding author Daniel Prieto-Alhambra, MD, Ph.D., of the University of Oxford, in the UK.

"To our knowledge, this is the first study to identify [cardiovascular disease risk factors](#) for osteoporotic individuals using data that is routinely collected and readily available."

More information: Marta Pineda-Moncusí et al, Estimating the incidence and key risk factors of cardiovascular disease in patients at high risk of imminent fracture using routinely collected real-world data from the UK, *Journal of Bone and Mineral Research* (2022). [DOI: 10.1002/jbmr.4648](#)

Provided by Wiley

Citation: Cardiovascular disease risk in adults with compromised bone health (2022, September 7) retrieved 29 January 2023 from <https://medicalxpress.com/news/2022-09-cardiovascular-disease-adults-compromised-bone.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--