


Study finds connection between symptomatic COVID-19 infection and impaired pain modulation

September 7 2022



Group differences in Condition Pain Modulation between symptomatic COVID-19, asymptomatic COVID-19, and control groups in the leg. *denotes significant difference between symptomatic and asymptomatic. IP = immediately post, 15P = 15minutes post. Credit: *The Journal of Pain* (2022). DOI: 10.1016/j.jpain.2022.06.010

A study led by researchers at the University of Oklahoma shows a connection between symptomatic COVID-19 infections and the ability of the central nervous system to respond to painful stimuli.

Christopher D. Black, Ph.D., associate professor in the Department of Health and Exercise Science, Dodge Family College of Arts and Sciences at OU, led the study with a doctoral student in his lab, Jessica A. Peterson. Peterson, the first author of the study, is now a postdoctoral researcher at the Pain Research and Intervention Center of Excellence at the University of Florida, Gainesville.

In late 2020, the [scientific community](#) began wondering what the potential long-term effects of COVID-19 infections might be. One of the common symptoms of COVID-19 [infection](#) is system-wide inflammation, which can lead to headaches and general aches and pains, also called myalgia.

"When you get inflammation, things that weren't painful before now become painful," Black said. "We realized we have all of these people who have had COVID-19, and there's a chance they may be at a higher risk of developing chronic pain conditions when they get older. We wanted to see whether people who had experienced a COVID infection—even though they no longer have symptoms—if they have normal function in their central nervous system in relation to their ability to 'correctly' respond to a painful stimuli."

The study, conducted from January through March 2021, included 59 participants separated into three groups. The symptomatic group included those who had previous symptomatic COVID-19 infections that were confirmed through PCR or antibody testing. An unexpected addition to the study became the asymptomatic group—those who had not known they had previously been infected by COVID-19 and had not experienced symptoms but were confirmed through antibody testing.

The control group had neither experienced COVID-19 infection nor had antibodies present at the time of testing.

The researchers measured each person's pain threshold—the amount of pressure applied to a muscle, in this case in the participant's arm or leg, that it takes for the participant to report feeling pain. Second, the researchers measured pain [modulation](#), or the signals sent from the central nervous system to the brain and spinal cord that can "change the volume" on how much pain a person reports.

To measure the pain modulation experienced by the test groups, the researchers followed the muscle pressure test with one of two experiments, a bout of knee extension exercise or a "cold presser test" where participants were asked to put their foot in a bucket of ice water for a brief period.

"These two tests, especially the ice water test (also called a conditioned pain modulation test), are used in [clinical settings](#) to help identify individuals who have or may be at a risk of developing a chronic pain condition," Black said. "So, we measured each person's pain threshold on their leg and on their arm. Then they would either do a bout of exercise with their leg and then we retest their pain threshold on the leg and the arm, or they would put their foot in the bucket of ice water and then we retest the leg and the arm. A 'normal' modulation response would result in an increase in their pain threshold—it would require greater pressure to be applied for them to tell us it hurt. However, in our symptomatic COVID participants, their pain thresholds did not change, indicating impaired conditioned pain modulation. Other studies have shown clear relationships between impaired pain modulation and the development of chronic pain conditions."

The study's finding shows a connection between participants who had experienced symptomatic, verified COVID-19 infections and the ability

of their central nervous system to respond normally to a conditioned pain modulation test.

"As far as we can tell, these are some of the first data on the effects of COVID-19 on pain outcomes," Black said. "We (in the U.S.) spend hundreds of billions of dollars on the treatment of [chronic pain conditions](#) every year. It's one of the leading causes of adult disability. Chronic pain is already a huge, chronic burden on the health care system, and we now have hundreds of millions of people in the U.S. who have had a COVID infection. The interaction of COVID and pain need to be further studied because if there is a chance of lasting impairments in pain modulation, this may put individuals at a greater risk in the future of developing a chronic [pain](#) condition."

The study was published July 21, 2022, in the *Journal of Pain*.

More information: Jessica A. Peterson et al, Symptomatic but not Asymptomatic COVID-19 Impairs Conditioned Pain Modulation in Young Adults, *The Journal of Pain* (2022). [DOI: 10.1016/j.jpain.2022.06.010](#)

Provided by University of Oklahoma

Citation: Study finds connection between symptomatic COVID-19 infection and impaired pain modulation (2022, September 7) retrieved 14 May 2023 from <https://medicalxpress.com/news/2022-09-symptomatic-covid-infection-impaired-pain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.