


Gel treats gum disease by fighting inflammation

September 20 2022



A topical gel that blocks the receptor for a metabolic byproduct called succinate treats gum disease by suppressing inflammation and changing the makeup of bacteria in the mouth. Credit: Yuqi Guo

A topical gel that blocks the receptor for a metabolic byproduct called succinate treats gum disease by suppressing inflammation and changing

the makeup of bacteria in the mouth, according to a new study led by researchers at NYU College of Dentistry and published in *Cell Reports*.

The research, conducted in mice and using [human cells](#) and plaque samples, lays the groundwork for a non-invasive treatment for [gum disease](#) that people could apply to the gums at home to prevent or treat gum disease.

Gum disease (also known as periodontitis or [periodontal disease](#)) is one of the most prevalent inflammatory diseases, [affecting nearly half of adults 30 and older](#). It is marked by three components: inflammation, an imbalance of unhealthy and [healthy bacteria](#) in the mouth, and destruction of the bones and structures that support the teeth. Uncontrolled gum disease can lead to [painful and bleeding gums](#), difficulty chewing, and tooth loss.

"No current treatment for gum disease simultaneously reduces inflammation, limits disruption to the oral microbiome, and prevents [bone loss](#). There is an urgent public health need for more targeted and effective treatments for this common disease," said Yuqi Guo, an associate research scientist in the Department of Molecular Pathobiology at NYU Dentistry and the study's co-first author.

Past research has linked increased succinate—a molecule produced during metabolism—to gum disease, with higher succinate levels associated with higher levels of inflammation. Guo and her colleagues at NYU Dentistry also [discovered in 2017](#) that elevated levels of succinate activate the succinate receptor and stimulate bone loss. These findings made the succinate receptor an appealing target for countering inflammation and bone loss—and potentially stopping gum disease in its tracks.

Strengthening the link between succinate and gum

disease

The researchers started by examining dental plaque samples from humans and blood samples from mice. Using metabolomic analyses, they found higher succinate levels in people and mice with gum disease compared to those with healthy gums, confirming what previous studies have found.

They also saw that the succinate receptor was expressed in human and mouse gums. To test the connection between the succinate receptor and the components of gum disease, they genetically altered mice to inactivate, or "knock out," the succinate receptor.

In "knockout" mice with gum disease, the researchers measured lower levels of inflammation in both the gum tissue and blood, as well as less bone loss. They also found different bacteria in their mouths: mice with gum disease had a greater imbalance of bacteria than did "knockout" mice.

This held true when the researchers administered extra succinate to both types of mice, which worsened gum disease in normal mice; however, "knockout" mice were protected against inflammation, increases in unhealthy bacteria, and bone loss.

"Mice without active succinate receptors were more resilient to disease," said Fangxi Xu, an assistant research scientist in the Department of Molecular Pathobiology at NYU Dentistry and the study's co-first author. "While we already knew that there was some connection between succinate and gum disease, we now have stronger evidence that elevated succinate and the succinate receptor are major drivers of the disease."

A novel treatment

To see if blocking the succinate receptor could ameliorate gum disease, the researchers developed a gel formulation of a small compound that targets the succinate receptor and prevents it from being activated. In laboratory studies of human gum cells, the compound reduced inflammation and processes that lead to bone loss.

The compound was then applied as a [topical gel](#) to the gums of mice with gum disease, which reduced local and systemic inflammation and bone loss in a matter of days. In one test, the researchers applied the gel to the gums of mice with gum disease every other day for four weeks, which cut their bone loss in half compared to [mice](#) who did not receive the gel.

Mice treated with the gel also had significant changes to the community of bacteria in their mouths. Notably, bacteria in the *Bacteroidetes* family—which include pathogens that are known to be dominant in gum disease—were depleted in those treated with the gel.

"We conducted additional tests to see if the compound itself acted as an antibiotic, and found that it does not directly affect the growth of bacteria. This suggests that the gel changes the community of bacteria through regulating inflammation," said Deepak Saxena, professor of molecular pathobiology at NYU Dentistry and the study's co-senior author.

The researchers are continuing to study the gel in animal models to find the appropriate dosage and timing for application, as well as determine any toxicity. Their long-term goal is to develop a gel and oral strip that can be used at home by people with or at risk for gum disease, as well as a stronger, slow-release formulation that dentists can apply to pockets that form in the gums during gum disease.

"Current treatments for severe gum disease can be invasive and painful.

In the case of antibiotics, which may help temporarily, they kill both good and bad bacteria, disrupting the oral microbiome. This new compound that blocks the succinate receptor has clear therapeutic value for treating gum disease using more targeted and convenient processes," said Xin Li, professor of molecular pathobiology at NYU Dentistry and the study's lead author.

Additional study authors include Scott Thomas, Yanli Zhang, Bidisha Paul, Sungpil Chae, Patty Li, Caleb Almeter, and Angela Kamer of NYU Dentistry; Satish Sakilam and Paramjit Arora of NYU Department of Chemistry; and Dana Graves of the University of Pennsylvania School of Dental Medicine.

More information: Xin Li, Targeting Succinate Receptor Effectively Inhibits Periodontitis, *Cell Reports* (2022). DOI: [10.1016/j.celrep.2022.111389](https://doi.org/10.1016/j.celrep.2022.111389). [www.cell.com/cell-reports/full ... 2211-1247\(22\)01226-8](https://www.cell.com/cell-reports/full...2211-1247(22)01226-8)

Provided by New York University

Citation: Gel treats gum disease by fighting inflammation (2022, September 20) retrieved 7 July 2023 from <https://medicalxpress.com/news/2022-09-gel-gum-disease-inflammation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--