

# Hair straightening chemicals associated with higher uterine cancer risk

17 October 2022



Credit: Pixabay/CC0 Public Domain

Women who used chemical hair straightening products were at higher risk for uterine cancer compared to women who did not report using these products, according to a new study from the National Institutes of Health. The researchers found no associations with uterine cancer for other hair products that the women reported using, including hair dyes, bleach, highlights, or perms.

The [study data](#) includes 33,497 U.S. women ages 35-74 participating in the Sister Study, a study led by the National Institute of Environmental Health Sciences (NIEHS), part of NIH, that seeks to identify [risk factors](#) for [breast cancer](#) and other [health conditions](#). The women were followed for almost 11 years and during that time 378 uterine cancer cases were diagnosed.

The researchers found that women who reported frequent use of hair straightening products, defined as more than four times in the previous year, were more than twice as likely to go on to develop uterine cancer compared to those who did not use the products.

"We estimated that 1.64% of women who never used hair straighteners would go on to develop uterine cancer by the age of 70; but for frequent users, that risk goes up to 4.05%," said Alexandra White, Ph.D., head of the NIEHS Environment and Cancer Epidemiology group and lead author on the new study. "This doubling rate is concerning. However, it is important to put this information into context—uterine cancer is a relatively rare type of cancer."

Uterine cancer accounts for about 3% of all new cancer cases but is the most common cancer of the female reproductive system, with 65,950 estimated new cases in 2022. Studies show that incidence rates of uterine cancer have been rising in the United States, particularly among Black women.

Approximately 60% of the participants who reported using straighteners in the previous year were self-identified Black women, according to the study published in the *Journal of the National Cancer Institute*. Although, the study did not find that the relationship between straightener use and uterine cancer incidence was different by race, the [adverse health effects](#) may be greater for Black women due to higher prevalence of use.

"Because Black women use hair straightening or relaxer products more frequently and tend to initiate use at earlier ages than other races and ethnicities, these findings may be even more relevant for them," said Che-Jung Chang, Ph.D., an author on the new study and a research fellow in the NIEHS Epidemiology Branch.

The findings are consistent with prior studies showing straighteners can increase the risk of hormone-related cancers in women.

The researchers did not collect information on brands or ingredients in the hair products the women used. However, in the paper they note that several chemicals that have been found in

straighteners (such as parabens, bisphenol A, metals, and formaldehyde) could be contributing to the increased uterine cancer risk observed. Chemical exposure from hair product use, especially straighteners, could be more concerning than other [personal care products](#) due to increased absorption through the scalp which may be exacerbated by burns and lesions caused by straighteners.

"To our knowledge this is the first epidemiologic study that examined the relationship between straightener use and uterine cancer," said White. "More research is needed to confirm these findings in different populations, to determine if hair products contribute to health disparities in [uterine cancer](#), and to identify the specific chemicals that may be increasing the risk of cancers in women."

This team previously found that permanent hair dye and straighteners may increase breast and ovarian cancer risk. That work was published in the *International Journal of Cancer*.

**More information:** Che-Jung Chang et al, Use of Straighteners and Other Hair Products and Incident Uterine Cancer, *JNCI: Journal of the National Cancer Institute* (2022). [DOI: 10.1093/jnci/djac165](https://doi.org/10.1093/jnci/djac165)

Carolyn E. Eberle et al, Hair dye and chemical straightener use and breast cancer risk in a large US population of black and white women, *International Journal of Cancer* (2019). [DOI: 10.1002/ijc.32738](https://doi.org/10.1002/ijc.32738)

The Sister Study:  
[sisterstudy.niehs.nih.gov/English/index1.htm](https://sisterstudy.niehs.nih.gov/English/index1.htm)

Provided by National Institutes of Health

APA citation: Hair straightening chemicals associated with higher uterine cancer risk (2022, October 17) retrieved 21 October 2022 from <https://medicalxpress.com/news/2022-10-hair-straightening-chemicals-higher-uterine.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*